

26th Annual General Pediatric Review & Self-Assessment


Pulmonology

Maria Franco, MD

Pediatric Pulmonology
Nicklaus Children's Hospital
Miami, FL

Nicklaus Children's Hospital logo

1



26th Annual General Pediatric Review & Self-Assessment

Disclosure of Relevant Relationship

Dr. Franco disclosed relevant conflicts of interests (COIs) and/or financial relationships in the past 24 months with the following ineligible companies:

Ineligible Company(ies)	Role/ Type of Relationship
Vertex Pharmaceutical	Research Support
PTC 124	Contracted Research

All COIs have been mitigated prior to this activity

Dr. Franco will support this presentation and clinical recommendations with the “best available evidence” from medical literature.

Dr. Franco does not intend to discuss an unapproved/investigative use of a commercial product/device in this presentation.

Nicklaus Children's Hospital logo

2

General Signs and Symptoms

3

Cough

- Acute:
 - Infection, aspiration, foreign body
- Recurrent
 - Reactive airways, CF, reflux, aspiration, anatomic abnormality, passive smoking
- Chronic (> 3 weeks)
 - RAD, CF, GER, Pertussis, anatomic abnormality, smoke exposure, psychogenic (often after a URI, does not occur at night)

4


Tachypnea

- WHO definition:
 - 0-2 mos > 60 bpm
 - 2-12 mos > 50 bpm
 - 12 months- 5 years of age > 40 bpm
 - > 20 bpm in older children and adolescents

5


Clubbing

- CF
- Bronchiolitis obliterans
- Primary Ciliary Dyskinesia
- Congenital heart disease
- Biliary atresia, IBD, alpha1antitrypsin
- Thyrotoxicosis, hypothyroid
- Idiopathic, hereditary



6

Sites & Sounds of Airway Obstruction



7

Definitions of obstructive noises

Stertor : respiratory sound characterized by heavy snoring or gasping

- Low-pitched
- Inspiratory
- Nasopharynx, oropharynx, nasal passage
- Loudest over neck, cheeks

(A heavy snoring inspiratory sound occurring in coma or deep sleep, sometimes due to obstruction of the larynx or upper airways)

8

Stertor (...Snoring)

Causes

- Choanal atresia
- Mandibular hypoplasia
- Macroglossia
- Nasal congestion
- Adenotonsillar hypertrophy
- Pharyngeal insufficiency
- Encephalocele
- Dermoid of base of tongue
- Thyroglossal duct cyst
- Lingual thyroid

9

Definitions of obstructive noises

STRIDOR : Harsh noise caused by turbulent flow

- Inspiratory = larynx
- Expiratory = trachea
- Biphasic = fixed lesion in subglottic region

10

Inspiratory Stridor

Causes

- Laryngomalacia
- Vocal cord paralysis
 - Unilateral – left (recurrent laryngeal nerve)
 - Bilateral – brainstem
- Laryngotracheoesophageal cleft
- Laryngocele
- Laryngeal polyp
- Abscess
- Hypotonia
- Vocal Cord Dysfunction


11

Inspiratory Stridor

Laryngomalacia

- 60% of insp. stridor
- 90% require no intervention
- Improves when prone (sleep)
- Worsens with activity and with presence of GERD
- Worsens over first 1-6mos, then improves
- Watch for FTT, apnea, cyanosis
- Bronchoscopy if does not follow classical history

12



Laryngomalacia


13

Biphasic Stridor

Causes


- Subglottic stenosis
 - Congenital
 - Acquired – intubation, croup
- Subglottic hemangioma
- Laryngeal web
- GERD

14




NORMAL LARYNX

15



SUBGLOTTIC STENOSIS

16



17

Biphasic Stridor

Causes

- **Croup** – acute barky cough, stridor, resp. distress
 - Low grade fever
 - Rhinorrhea
 - Worse at night
 - 3mos-3yrs
 - Parinfluenza 1-3, RSV, Influenza
 - Fall/Winter

18

Biphasic Stridor

Causes

- **Recurrent Croup**
 - Consider underlying airway anomaly
 - GERD
 - Spasmodic (reactive airways)

19

Definitions of obstructive noises

Wheeze : Higher-pitched expiratory noise

- Monophonic, homophonous = large airway
= **expiratory stridor**
- Polyphonic, heterophonous, musical = small airways

20

Homophonous Wheezing – Expiratory Stridor

Causes


- Tracheobronchomalacia
 - Deficient cartilage rings
 - Worse with exertion, agitation
 - Prolonged expiratory phase
 - Narrow trachea on expiratory lateral films
 - Primary vs. Secondary
 - BPD, TEF, vascular anomalies

21

Vascular Rings

- Tracheal or tracheobronchial malacia and stenosis may develop in association with some of these lesions in the areas where the greatest degree of compression exists. This is particularly true in cases of anomalous left pulmonary artery.


22



Aberrant left pulmonary artery or pulmonary artery sling.


Most patients are symptomatic by 1 month after birth.

23



24


Double aortic arch



The double aortic arch forms a ring around the trachea and esophagus, compressing both of these structures. Tracheoesophageal compression typically results in early symptoms

25

Double Aortic Arch



26

Mediastinal Mass

27

Bronchogenic Cyst
In a 12 month old
Evaluated for chronic
wheezing



28

Heterophonous Wheezing

Causes

- Asthma
- Bronchiolitis
- Pneumonia (Mycoplasma)
- GERD – inflammation, bronchospasm
- Heart Failure – often presents around 2 mos
- Cystic Fibrosis
- Ciliary Dyskinesia
- Food Allergy

29

ASTHMA

30

Asthmatic Inflammation

Asthma Is a Chronic Inflammatory Disease

- Asthma is a chronic inflammatory disease.
- Airway inflammation leads to:
 - Hyperresponsiveness—responses to triggers
 - Obstruction—usually reversible
 - Symptoms—cough, wheezing, dyspnea

NAEPP. *Clinical Practice Guidelines: Expert Panel Report 2: Guidelines for the Diagnosis and Management of Asthma*. June 2002. NIH publication 02-4051.

31

ASTHMA

Most common chronic
pediatric disorder

32

ATOPIC WHEEZING ASTHMA

- **MORE THAN ½ OF ALL CASES OF PERSISTENT ASTHMA START BEFORE AGE 3.**
- **80% START BEFORE 6 YEARS OF AGE**

33

ASTHMA DIAGNOSIS

Consider asthma if:

- History of chronic dry cough
- Cough during exercise
- Wheezing
- Wheezing with upper respiratory infections and or with exercise.

34

GOALS OF ASTHMA THERAPY

- Lack of symptoms
- Diminished response to triggers
- Full participation in usual activities
- Normal pulmonary function
- Use of as little medication as possible but as much as necessary

35

Pharmacologic Therapy

Quick-relief medications: ACUTE

- Short-acting Beta2-agonists:
Bronchodilatation
Albuterol
Levalbuterol
- Oral Corticosteroids : Anti-inflammatory
Up to 10 days of therapy no need for tapering dose.
- Ipratropium Bromide : Add on severe asthma

36

Pharmacologic Therapy

LONG TERM CONTROL MEDICATION

- Cromolyn Sodium
- Nedocromil
- Leukotriene Modifiers
- Long Acting Beta2-agonists
- Sustained-released theophylline
- Oral Steroids
- Inhaled Steroids
- Inhaled Steroids + Long Acting Beta2-agonists
- Biologics

37

- All medications decrease airway hyperactivity
- Cromolyn, Nedocromyl, Steroids and Leukotriene Modifiers also decrease inflammatory component of the airway.

38

ICS Safety Includes ...

- Data demonstrating no effect on final adult height at recommended doses
- Minimizing effects of chronic inflammation in the lung
- Protection of lung function
- Reducing risk of death, hospitalizations, and emergency room visits for asthma
- Significant reduction in asthma symptoms and exacerbations



39

“RULES OF TWO™”*

Patients Are Candidates for Maintenance Therapy If ...

- They are using a quick-relief inhaler more than 2 times per week
- They awaken at night due to asthma more than 2 times per month
- They refill a quick-relief inhaler prescription more than 2 times per year

*“RULES OF TWO™” is a trademark of the Baylor Health Care System.

40

Rule of Twos

- ≥ 2 days/week but not daily, ≥ 2 nights/month
→ **Mild persistent**
 - Treatment: **Low-dose inhaled corticosteroid
Leukotriene Modifiers**
- Daily daytime symptoms, > 1 night/week →
 - Treatment: **Moderate persistent**
Low-dose ICS + LABA, OR medium-dose ICS
- Continual symptoms → **Severe persistent**
 - Treatment: **High-dose ICS + LABA, and OCS
Or Biologics (monoclonal antibodies)**

41

Exercise can trigger asthma

- Symptoms are worse with cold, dry air
- However, exercise helps lungs function better and prevents obesity
- As long as asthma is well-controlled and a short-acting bronchodilator (rescue medicine) is used beforehand, children with asthma should be able to do sports
- Pulmonary function testing best first test; then exercise testing.

42

Bronchiolitis

43

Bronchiolitis

- Most common cause is RSV.
- Most common cause of first episode of wheezing in infants.
- Diagnosed by RSV antigen by nasal wash.
- For treatment steroids or bronchodilators are not recommended.
- Observation and close monitoring during hospitalization. Hydration when necessary.
- Nasal Suction prior feedings to avoid aspiration.

44

Acute Respiratory Failure

- **Increased Respiratory Drive**
 - Tachypnea – increased RR
 - Dyspnea – breathlessness
 - Retractions
 - Accessory muscle use
- **Decreased Respiratory Drive**
 - Decreased RR
 - Lethargy
 - Confusion
 - Snoring

45

Increase CO₂ is always the most clear sign of respiratory failure

Normal CO₂ is from 40 to 45 mm Hg

46

Acute Respiratory Failure

- Respiratory Muscle Fatigue
 - Paradoxical “see-saw” respirations
 - Grunting
 - Uncoordinated breathing
- Hypoxemia – $\text{PaO}_2 < 60$
- Hypercapnea – $\text{PaCO}_2 > 50$

47

Blood Gases

pH	PCO ₂	PO ₂	HCO ₃	
7.24	60	50	24	ARF
7.35	60	50	34	CRF
7.40	40	70	24	Normal

48

CYSTIC FIBROSIS

- Chronic, progressive and life limiting **autosomal recessive** genetic disease characterized by chronic respiratory disease, pancreatic insufficiency, elevation of sweat electrolytes and male infertility




49


CF

- CF is caused by a mutation in a gene located in the long arm of chromosome # 7.
- Gene called **CFTR** (**c**ystic **f**ibrosis **t**ransmembrane conductance **r**egulator)
- Membrane protein, epithelial chloride channel


50



51



53



55


5 Classes of CFTR Mutations

The diagram shows five classes of CFTR mutations, each represented by a cross-section of an airway cell. Class I shows a defective protein (red X). Class II shows a defective protein (red X). Class III shows a defective protein (red X). Class IV shows a defective protein (red X). Class V shows a reduced amount of protein (smaller blue structure).

I	II	III	IV	V
Defective Protein Production	Defective Protein Processing	Defective Protein Regulation	Defective Protein Conductance	Reduced Amounts CFTR protein

CF Mutations can be classified by the effect they have on the CFTR protein.

56



57

CF mutations

- There about 1700 known mutations of the CF gene
- The most common Delta F 508 (a type II mutation)

58

CF: Genetics

- More than 80 percent of patients are diagnosed by age three; however, nearly 10 percent of newly diagnosed cases are age 18 or older.
- Newborn screening Only Detects 10% of the cases

59

Presentation (CF PANCREAS)

- C** Chronic respiratory disease
- F** Failure to thrive
- P** Polyps
- A** Alkalosis, metabolic
- N** Neonatal intestinal obstruction
- C** Clubbing of fingers
- R** Rectal prolapse
- E** Electrolyte ↑ in sweat
- A** Aspermia / absent vas deferens
- S** Sputum – S.aureus/P.aeruginosa

60

CF: Symptoms

- Chronic sino-pulmonary disease
 - Chronic Sinusitis
 - Nasal Polyps
- Gastrointestinal/nutritional problems
- Salt-loss syndromes
- Normal intellect
- Life span about 30+ years
- If pancreatic sufficient, life span 56 years

61

SYMPTOMS

- People with CF have a variety of symptoms including:
 - Very salty-tasting skin
 - Persistent coughing, at times with phlegm; wheezing or shortness of breath
 - Excessive appetite but poor weight gain; and greasy, bulky stools.
 - Symptoms vary from person to person

62

CF: Gastrointestinal Disease

- Pancreatic insufficiency/malabsorption
- Neonatal intestinal obstruction Meconium Ileus(15%)
- Lipo-soluble vitamin deficiency
- Failure to thrive
- Recurrent distal intestinal obstruction
- Biliary stasis

63

CF: Pancreas-malabsorption




64

DIAGNOSIS



65

The sweat test (Chloride)



Negative (02/2017)

Under 30 mEq/L
Past was less than 40

Borderline

30-59 mEq/L

Positive

Over 60 mEq/L

[J.Peds Volume 181, Supplement](#), February 2017, Pages S4-S15.e1

66



The image displays the components of the MACRODUCT SYSTEM. It includes a black plastic device with a white sensor, a photograph of a newborn baby with a blue sensor on their chest, a white electronic control unit with various ports, and a red and white sensor strip. The text 'MACRODUCT SYSTEM' is prominently displayed in white on a blue background.

MACRODUCT SYSTEM

Start at 2 weeks of age

67

Blood Gases Patient with CF

- **Ph Metabolic Alkalosis**
 - **Low Sodium**
 - Normal Potassium
 - **Low Chloride**
 - **High Bicarbonate**
 - Normal BUN

68

One frequent differential Diagnosis

■ Ciliary Dyskenesia Primary:

- Recurrent Otitis Media
- Chronic Sinusitis
- Productive cough
- Bronchiectasis
- Recurrent Pneumonia
- Situs Inversus (Kartagener's Syndrome)

69

Differential Diagnosis for Wheezing

FOREIGN BODY ASPIRATION

70

Upper Airway Laryngeal area

- **Acute episode of: +/-**
 - Hoarse voice**
 - Cough**
 - Wheezing**
 - Respiratory distress**
 - No fever**
 - With or without history of choking episode.**

71

Airway Symptoms Lower

- **Persistent paroxysmal cough after choking episode.**
- **Localized wheezing does not improve with bronchodilators.**
- **Chest x-ray shows one side hyperinflation during inspiration and expiration from the side where the foreign body is located.**
- **Treatment Bronchoscopy.**

72

Pneumothorax

- **Primary spontaneous** – Teens and young adults, esp. tall thin males, Ehlers-Danlos, and Marfan.
- **Secondary spontaneous** – underlying conditions: pneumonia, empyema, cyst, Foreign body, asthma. Not due to trauma.
- **Iatrogenic**
- **Catamenial pneumothorax** – associated with menses, passage of intra-abdominal air through diaphragmatic defects.

73

PNEUMOTHORAX

- **ACUTE ONSET OF CHEST PAIN, RESPIRATORY DISTRESS IN A PATIENT WITH ASTHMA**
- **TRAUMA**
- **ACUTE DECOMPENSATION IN A PATIENT ON MECHANICAL VENTILATION**

ALWAYS DECREASE BREATH SOUNDS IN THE LUNG WITH THE PNEUMOTHORAX

74

Pneumothorax

- Rx: <5% is “small”, may resolve spontaneously, give 100%O₂,
- >20% = “large” pneumos need chest tube
- Tension pneumos need immediate decompression.

75

Hemoptysis

- ◆ The leading causes of hemoptysis in children are cystic fibrosis, congenital heart disease (CHD), and trauma.
- ◆ Infection, tracheostomy-related, bronchiectasis, foreign body, A-V Malformation, trauma, tumor.
- ◆ Pulmonary hemosiderosis, Goodpasture syndrome, Wegener granuloma

76


Epistaxis

- ◆ Most Common Cause is: **digital trauma**
- ◆ Adolescents – question about drug abuse.
- ◆ Kiesselbach plexus
- ◆ Order coagulation and hematologic studies only if prolonged, severe, or family history.

77

PULMONARY FUNCTION TEST

78



79


Obstructive Disorders

Volume (L)

- Characterized by a limitation of expiratory airflow
 - Examples: Asthma, COPD
- **Decreased:** FEV₁, FEV₂₅₋₇₅, FEV₁/FVC ratio (<0.8)
- **Increased or Normal:** TLC
- Normal Diffusion Capacity

80


Restrictive Lung Disease



- **Characterized by diminished lung volume due to:**
 - change in alteration in lung parenchyma (interstitial lung disease)
 - disease of pleura, chest wall (e.g. scoliosis), or neuromuscular apparatus (e.g. muscular dystrophy)
- **Decreased TLC, FVC**
- **Normal or increased: FEV₁/FVC ratio**
- **Diffusion capacity could be altered**

81


Large Airway Obstruction



- **Characterized by a truncated inspiratory or expiratory loop**

82

Spirometry Loops



83

Thank you !!!



GOOD LUCK!!!

84