

The Annual General Pediatric Review & Self Assessment

# CARDIOLOGY

**Jason Katz, MD**

Attending, Pediatric Cardiology  
Director, Pediatric Cardiology Fellowship Program  
Nicklaus Children's Pediatric Specialists  
Nicklaus Children's Hospital  
Miami, Florida

## The Annual General Pediatric Review & Self Assessment

# Disclosure of Relevant Relationship

Dr. Katz has not had (in the past 24 months) any relevant conflicts of interest or relevant financial relationship with the manufacturers of products or services that will be discussed in this CME activity or in his presentation.

Dr. Katz will support this presentation and clinical recommendations with the “best available evidence” from medical literature.

- Dr. Katz does not intend to discuss an unapproved/investigative use of a commercial product/device in this presentation.

# Passing the Boards

- High Yield Review
  - Presentations
    - Murmurs
    - CHF
    - Cyanosis
    - Shock
  - Syndromes
  - Scenarios
  - EKGs
  - Arrhythmias

# Presentation of CHD

- Murmur
- CHF
- Cyanosis
- Shock
- High Risk of Heart Disease
  - Down Syndrome, Trisomy 13, 18, etc..

# Murmurs

- Ejection Systolic Murmur
  - Pulmonary stenosis and TOF (LUSB)
  - Aortic stenosis (RUSB)
- Holosystolic Murmurs
  - VSD (LLSB) and Mitral regurgitation (apex)
- Continuous Murmurs
  - PDA (machine-like) (LUSB - left infraclavicular)
  - BT Shunt murmurs (throughout chest)
- Innocent Murmurs
  - PPS (Peripheral pulmonic stenosis) classically heard in infants
    - Ejection systolic (ULSB) radiating to both axilla
  - Still's murmur (classically 3 to 6 year old) (LSB to apex)
    - Vibratory sounding - louder in supine position
  - Venous hum (disappears when the head is turned to the side)
    - Supra and infra clavicular areas

# CHF

- Congestive Heart Failure
  - From increased pulmonary blood flow
    - VSD, PDA, TGA, HLHS, Truncus Arteriosus
  - Decrease ventricular function
    - Dilated cardiomyopathy
  - Signs and Symptoms
    - Tachypnea (classic sign is comfortable tachypnea)
    - Tachycardia
    - Poor po intake / tires with feeds (sweats w/ feeds)
    - Poor weight gain

# Cyanotic baby work-up


- Increased pulmonary vascular markings (white lungs on CXR)
  - TGA (increased or normal) – comfortable tachypnea
  - TAPVR – respiratory distress
  - HLHS – shock (mottled, acidotic, pulseless)
- Decreased pulmonary vascular markings (Dark lungs on CXR)
  - Tricuspid Atresia – Left superior axis on EKG
  - Pulmonary Atresia
  - Critical Pulmonary Stenosis and TOF – harsh SEM

# Shock

- HLHS (Hypoplastic Left Heart Syndrome)
  - Cyanotic (often with sats in high 80's/low 90's)
  - Can present in shock (ductal dependent Systemic BF)
  - Start prostin ASAP
- Critical Coarctation & Critical Aortic Stenosis
  - Can present in shock (ductal dependent Systemic BF)
  - Start prostin ASAP



# Critical Coarctation



# Prostin

- Start a prostin infusion to keep ductus open
  - If suspect ductal dependent lesion (Echo)
    - Any neonate who presents in shock
    - Any neonate with severe hypoxemia
- Dose
  - To open ductus after closure
    - 0.1 mcg / kg / min
  - To keep ductus open (can use for weeks)
    - 0.02 - 0.05 mcg / kg / min
- Complications (decreased by lower doses)
  - Apnea, hypotension, fever

# Diagnostic Tests

- History & Physical
- Oxygen Saturation
- 4 extremity BPs
- Hyperoxia test
- EKG
- Echo
- (Cardiac Cath, EP Study, Holter)

# Treatment

- MURMUR
  - Depends on heart defect
- CHF
  - Lasix, Lasix, Lasix, Digoxin, and good nutrition
- HYPOXEMIA
  - Nothing or if ductal dependent than Prostin infusion
- SHOCK
  - Start Prostin infusion, inotropes and fluid resuscitation


# Treatment

- Ultimately most patients with congenital defects require :
  - Surgical correction
    - i.e. VSD closure, TGA repair (ASO), TOF repair
  - Surgical palliation
    - i.e. PA band, Norwood, Glenn, Fontan


# Acyanotic Heart Defects

- PDA
- Coartation of the Aorta
- ASD - soft ESM from relative PS and fixed split S2
- VSD
- AV Canal

# Patent Ductus Arteriosus (PDA)




# VSD Physiology






# AV Canal Defect



# Superior Axis




# Cyanotic Heart Defects

- Tricuspid Atresia
  - Cyanosis, maybe ductal dependent PBF
- Pulmonary Atresia & Critical pulmonary Stenosis
  - Cyanosis, probable ductal dependent PBF
- TAPVR (Total Anomalous Pulmonary Venous Return)
  - Cyanotic, tachypneic, comfortable
  - Obstructed type (very cyanotic and sick)
    - Surgical emergency

# Cyanotic Heart Defects

- TGA (Transposition of the Great Arteries)
  - Cyanosis, tachypneic but comfortable, no murmur
  - Most common cyanotic lesion presenting at birth
    - May need balloon atrial septostomy (if very blue)
    - Surgical repair as neonate

# Transposition of The Great Arteries




# D-TGA (Egg on a string)



# Cyanotic Heart Defects

- TOF (Tetrology of Fallot)
  - Ejection Systolic Murmur from birth and cyanosis
  - May present at birth or weeks later
  - Full surgical repair usually at 4 - 6 months oldSometimes full repair as neonate or just a BT shunt

# Tetralogy of Fallot



- 1- Pulmonary Stenosis
- 2 - Right Ventricular Hypertrophy
- 3 - Overriding Aorta
- 4 - VSD



# TOF (Boot shaped Heart)




# HLHS


- HLHS (Hypoplastic Left Heart Syndrome)
  - Cyanotic (often with sats in high 80's/low 90's)
  - Usually tachypneic
  - Can present in shock (ductal dependent Systemic BF)
  - Start prostin ASAP
  - Presents as neonate
  - 3 stage surgical palliation
    - Norwood - 1<sup>st</sup> week of life
    - Bidirectional Glenn - 4 months old
    - Fontan - 2 to 4 years old

# Normal

# HLHS




# BT shunt vs Sano




# BDG


**Bi-directional Glenn Shunt**  
Temporary surgical procedure as an intermediate  
procedure for hearts with only one usable ventricle  
(In this illustration Tricuspid Atresia)




# Extracardiac fontan



# Superior Axis (blue with a murmur)




# Tricuspid Atresia





# Down Syndrome


(Atrioventricular Septal Defect)



# Syndromes

- Down Syndrome
  - VSD, Atrioventricular septal defect
  - Pink, holosystolic murmur
  - Superior axis on EKG (negative in lead AVF)

# Superior Axis (Pink with murmur)



# Williams Syndrome

Supravalvar Aortic Stenosis



# Syndromes

- William Syndrome
  - Supravalvar Aortic Stenosis
  - Pink, ejection systolic murmur
  - Autosomal Dominant

# Turner's Syndrome

Coarctation of the Aorta and Aortic Stenosis



[medgen.genetics.utah.edu](http://medgen.genetics.utah.edu)

# DiGeorge Syndrome (VCF/22q1.1)

TOF/IAA/Truncus (conotruncal defects)



# Charge Syndrome





# Noonan's Syndrome

HCM, PS, ASD



# Syndromes


- Turner's Syndrome - Coarctation of the Aorta, AS
- DiGeorge Syndrome/22q1.1 del - TOF, IAA, Truncus
- CHARGE Syndrome - Wide variety
- Noonan Syndrome - HCM, PS, ASD (Autosomal Dominant)

# Other Genetic diseases

- Tuberos Sclerosis - cardiac rhabdomyomas
- Marfans - aortic root aneurysm, MVP (AD)
- Romano-Ward - LQTS (AD)
- Jervell & Lange-Nielsen - LQTS & Deafness (AR)

# Scenarios

- LQTS (Long QT syndrome)
  - QTc > 460ms (QT / square root of previous RR)
  - Alarm clock, swimming pool
  - Common treatment is Beta blocker
  - May need automatic internal cardiac defibrillator (AICD)
    - Refer to pediatric cardiology
- Maternal Diabetes
  - Neonate with hypertrophic cardiomyopathy
  - Usually gets better with time
- Fetal Echo
  - Can miss VSDs, Coarctation of the Aorta



# Rheumatic Fever

- History of recent group A strep pharyngitis
  - Polyarthrititis
  - New Murmur (MR, AR)
  - Sydenham's chorea
  - Erythema marginatum
  - Subcutaneous nodules
- Treatment
  - Aspirin for arthritis relief and Penicillin course
  - Penicillin for prophalaxis to prevent repeat infection


# Kawasaki Disease

- Need fever x 5 days or more and 4 out of 5 criteria below:
  - Eyes - Non-exudative conjunctivitis
  - Lips /mouth - erythema, cracked lips or strawberry tongue
  - Neck - unilateral cervical lymphadenopathy
  - Hands and feet - edema/erythema - eventually peeling
  - skin
- Treat with ASA and IVIG
  - Decreases risk of coronary artery anerysms
  - Echo to look at coronaries and ventricular function
    - Initial echo for baseline and rpt in 2 - 3 wks

# Chest Pain

- Benign if:
  - Reproducible to palpation
  - Worse with deep breathing
- Red flags
  - happens with exercise!!!!
  - If yes - cardiology referral and not allowed to exercise
- If worse when lying down think pericarditis
  - Friction rub and possible history of recent URI





**Pericarditis** Electrocardiogram in acute pericarditis showing diffuse up-sloping ST segment elevations seen best here in leads II, III, aVF, and V3 to V6. There is also subtle PR segment deviation (positive in aVR, negative in most other leads). ST segment elevation is due to a ventricular current of injury associated with epicardial inflammation; similarly, the PR segment changes are due to an atrial current of injury which, in pericarditis, typically displaces the PR segment upright in lead aVR and downward in most other leads. Courtesy of Ary Goldberger, MD.


# Syncope

- Vasovagal Syncope (benign)
  - Caused by pain, fear, standing too long
  - Treat with increasing salt and fluids in diet
- Syncope (malignant)
  - Happens with exercise
    - Coronary abnormality
    - Hypertrophic cardiomyopathy
  - Fast onset (arrhythmia) - patient usually falls and gets hurt
  - Long QT syndrome
  - Refer for cardiology consult (no exercise until evaluation)


# EKGs

- Sinus arrhythmia - due to respiratory variation
- Premature atrial contraction (PAC or APB)
- Premature ventricular contraction (PVC or VPB)


# Sinus Arrhythmia



HEALTH INTERACTIVE © 1999 - WWW.RNCEUS.COM




**Atrial premature beats** The third beat is an atrial premature beat that is normally conducted. The interval between the second sinus beat and the ectopic beat is shorter than the interval between the first two sinus beats. The P wave morphology differs from that of sinus rhythm. Since the RR cycle length is shorter, there is a decrease in the rate of conduction of the ectopic beat through the atrioventricular node. The PR interval is therefore longer than that of the sinus beat. Activation of the ventricular myocardium occurs in a normal fashion; as a result, the QRS complex is unchanged from that of sinus rhythm.



**Ventricular premature beat** The fourth beat is a ventricular premature beat (VPB). It has a wide, bizarre morphology, with a duration  $>0.16$  seconds.


# EKGs

- AV Block (1<sup>st</sup> 2<sup>nd</sup> and 3<sup>rd</sup> degree)




**First degree atrioventricular (AV) block** First degree AV block is caused by a prolongation or delay in impulse conduction through the AV node. It is defined as a PR interval  $>0.20$  seconds. In this case the PR interval (blue lines) is approximately 0.22 seconds.





**Mobitz type I second degree atrioventricular (AV) block** Several characteristic features of Wenckebach second degree AV block are noted on this ECG (P waves are marked by arrows): 1) The PR interval after the second P wave is longer than the preceding PR interval; the third P wave is not conducted at all. 2) The PP intervals are constant; however, the RR interval surrounding the completely blocked P wave is longer than the preceding normal RR interval. 3) The fourth P wave (after the blocked beat) is conducted normally; this PR interval is shorter than the PR interval that immediately preceded the pause.




**Mobitz type II second degree atrioventricular (AV) block** The third and sixth P waves are not conducted through the AV node (there is no associated QRS complex). The PR interval is constant prior to and after the non-conducted beats.



**Third degree atrioventricular block** The P waves are completely dissociated from the QRS complexes. The QRS complexes are narrow, indicating a junctional escape rhythm. The atrial and ventricular rates are stable; the former is faster than the latter.

# EKGs


- WPW
  - Delta wave (pre-excitation)
  - Short PR
  - SVT
  - Treat Beta-blocker not with Digoxin



**Wolff-Parkinson-White syndrome** The two main electrocardiographic features of Wolff-Parkinson-White syndrome (WPW) include a short PR interval ( $<0.12$  seconds) and a delta wave (red arrows). The QRS complex is wide ( $>0.12$  seconds) and represents a fusion beat; the initial portion (delta wave) results from slow ventricular activation via the accessory pathway, while the termination of ventricular activation is via the normal conduction system leading to a fairly normal terminal portion of the QRS.

# Atrial Tachyarrhythmias

- Re-entrant Tachycardia
  - Fast on / Fast off
    - Abrupt onset / termination
  - AVRT, AVNRT, AFlutter
- Ectopic Tachycardias
  - Warm up and cool off
    - Gradually gets faster
    - Gradually gets slower
  - JET, EAT, Sinus Tach



# Supra Ventricular Tachycardia

## Reentrant tachycardia


- Narrow complex, fast (>200), not variable
- Abruptly starts and abruptly stops
- Treatment
  - Vagal maneuver – ice, temperature, PIV
  - Adenosine IV given as a **push** – very fast!!!
    - Very short half-life, so must push fast
    - Preferable in upper extremity
    - Can cause VFib

# SVT: Presentation

- Infants
  - Poor feeding
  - Pallor
  - S+S of CHF
- Older Children
  - Palpitations
  - Chest pain/discomfort




SVT HR > 200 in children




**AV nodal reentrant tachycardia** Junctional tachycardia or atrioventricular nodal reentrant tachycardia is a supraventricular tachyarrhythmia that originates within the atrioventricular node. The rate is generally between 140 to 220 beats per minute and there is usually a 1:1 atrial-ventricular association; as a result, every QRS complex has an associated P wave. In this case, atrial and ventricular activation are simultaneous; as a result, the P wave is superimposed upon the QRS complex

# Ventricular Arrhythmias


- Pulseless Rhythms
  - Too fast to fill the heart and maintain a BP
  - Ventricular Tachycardia
  - Ventricular Fibrillation
  
- Treatment
  - Asynchronous Defibrillation



**Nonsustained ventricular tachycardia** Nonsustained ventricular tachycardia in a patient with underlying atrial fibrillation. The ventricular arrhythmia consists of nine beats at an approximate rate of 170 beats/min. Courtesy of Ary Goldberger, MD.




**Monomorphic ventricular tachycardia** Three or more successive ventricular beats are defined as ventricular tachycardia (VT). This VT is monomorphic since all of the QRS complexes have an identical appearance. Although the P waves are not distinct, they can be seen altering the QRS complex and ST-T waves in an irregular fashion, indicating the absence of a relationship between the P waves and the QRS complexes, i.e. AV dissociation is present.



**Polymorphic ventricular tachycardia** The QRS complexes have markedly differing morphologies due to changes in the direction (vector) of myocardial activation.

V<sub>2</sub>



**Ventricular fibrillation** There is a complete absence of properly formed QRS complexes and no obvious P waves. A recent onset (eg, within minutes) of the arrhythmia is suggested by the coarse morphology of the fibrillatory waves.

# Antiarrhythmic drug classification

- Class I - Na channel blocker
  - Class IA: Quinidine, Procainamide
  - Class IB: **Lidocaine**
  - Class IC: Propafenone, Flecainide
- Class II-Beta-blockers (propranolol, esmolol)
- Class III-K channel blocker (**Amiodarone**)
  - Adverse reactions of Amiodarone
    - Hypotension, Torsade, QT prolongation, Severe bradycardia, AV block
    - Hypo/hyperthyroidism, pulmonary toxicity, blue-grey skin discoloration
- Class IV-Ca channel blocker (Verapamil, Diltiazem)
  - Do not use if under one year old

# Some Tips

- Stay away from answers with **Always** and **Never**
- Referring patient is usually not a good answer
- Start with the most basic test or study
- They love to give you normal patients that just need reassurance (always consider this as an option)



# Thank you

- Good Luck
- Any questions contact me at [jason.katz@mch.com](mailto:jason.katz@mch.com)

# Question #1

- 13 year male seen in your office for a history of chest pain. He explains the chest pain is located in the left side of his chest, no radiation and resolves spontaneously. It worsens with deep breathing and is reproducible on your exam. He also explains his grandfather had a myocardial infarction at the age of 62. Of the following, the MOST appropriate approach for this patient at this time is
  - A. Cardiac catheterization
  - B. Reassurance with clinical follow-up
  - C. Echocardiography
  - D. Electrocardiogram

## Question #2

- Newborn baby born full term via NSVD who at 12 hours of age is noted to be mottled and tachypneic. Upon arrival in the NICU, his arterial blood gas shows a bicarbonate of 12 and a  $po_2$  of 42. Of the following, the best management plan is
- A. Order an echocardiogram
- B. Start a prostin infusion
- C. Give a 10cc/kg bolus of normal saline
- D. Start an infusion of epinephrine

# Question #3

- A two month old is seen in your office for a well baby visit. You note he has a loud holosystolic murmur heard best at the left midsternal border. His mom states he is now taking longer to finish his feeds and seems to be sweating and breathing harder with his feeds. He has gained no weight over the past month. Of the following, the best management plan is
- A. Start propranolol
- B. Admit to the hospital for immediate surgery
- C. Reassure mother that baby is OK
- D. Start furosemide

# Question #4

- A six month old male seen in the ER for a history of poor feeding for the past 24 hours. The baby appears to be tachypneic and pale. Once the baby is placed on a telemetry monitor it is noted the baby's heart rate is 240bpm. The blood pressure and oxygen saturation are normal. Of the following, the best management plan is
  - A. Start furosemide
  - B. Synchronized cardioversion
  - C. Place an IV and give adenosine rapidly
  - D. Start propranolol

# Question #5

- A 14 year old girl is found unconscious in a swimming pool after diving into the pool. She is taken out and found to be in pulseless ventricular fibrillation. She is shocked back into normal sinus rhythm and is brought to the hospital in good condition. Of the following, the appropriate diagnostic study to start with is
  - A. Echocardiogram
  - B. Electrocardiogram
  - C. Chest x-ray
  - D. Cardiac catheterization